AUNCLASSIFIED A GOOD TO SIFIED Armed Services Technical Information Higency

Reproduced by DOCUMENT SERVICE CENTER KNOTT BUILDING, DAYTON, 2, 0 HIO

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address: Armed Services Technical Information Agency, Document Service Center, Knott Building, Dayton 2, Ohio.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U.S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

Best Available Copy

Project RANDUM

This is a working paper. It may be expanded, modified, or withdrawn at any time. The views, conclusions, and recommendations expressed herein do not necessarily reflect the official views or policies of the U.S. Air Force.

-7he RAND Corporation

SANTA MONICA . CALIFORNIA-

PROJECT RAND

RESEARCH MEMORANDUM

Notes on Linear Programming: Part XV
MINIMIZING THE NUMBER OF CARRIERS
TO MEET A FIXED SCHEDULE

by

G. B. Dantzig D. R. Fulkerson

RM-1328

24 August 1954

Assigned to

This is a working paper. It may be expanded, modified, or withdrawn at any time. The views, conclusions, and recammendations expressed herein da nat necessarily reflect the official views or policies of the United States Air Force.

Authority to Reproduce

Granted to ASTIA-DSC

Per Rough 14r 19 axie 56

The

RHIII) Corporation

1700 MAIN ST. . SANTA MONICA . CALIFORNIA

Copyright, 1954
The RAND Corporation

SUMMARY

It is shown that the problem of determining the minimum number of carriers required to meet a fixed schedule of transportation can be made into a linear programming problem.

MINIMIZING THE NUMBER OF CARRIERS TO MEET A FIXED SCHEDULE

by

G. B. Dantzig
D. R. Fulkerson

1. INTRODUCTION

C. Tompkins (see [3]) has given a discrete idealization of a scheduling problem which arose in the routing of Navy fuel oil tankers. A combinatorial problem of this kind has also been discussed by J. Robinson and J. Walsh [2], together with a proposed method of computation. The algorithm they outline, however, fails to narrow the number of possibilities sufficiently to make it a feasible computational method for most problems.

In this note we show how the tanker scheduling problem can be made into a linear programming problem of transportation type, albeit large. The size of the system is mitigated somewhat by the following facts: (1) most of the variables are constrained to be zero, (2) the minimizing form is particularly simple, and (3) even a large transportation type problem having no special features can be solved by hand using the simplex algorithm [1].

2. THE PROBLEM

A rectangular array of spaces is furnished, one row for each pickup point and one column for each discharge point. In each space (i,j), $i=1,2,\ldots,m$; $j=1,2,\ldots,n$ is a sequence of numbers $t_{i,j}^k$, $k=1,2,\ldots$ representing the times at which a tanker is to load fully at pickup point i to deliver to destination j. For example, in the array

			1				2	2			3	
1	1,	4,	7,	10,	13		9,	15		É	5, 12	2
2	3,	6,	9,	12		7,	10,	13,	15	5,	10,	15

the sequence 3, 6, 9, 12 in box (1,2) means that at these times a tanker is to begin loading at pickup point 2 for delivery to discharge point 1. Multiple loads to go from i to j can be taken care of by repetitions in the sequence $t_{i,j}^k$. We make the further assumption, not made in [3], that the total number of entries in the table $t = (t_{i,j}^k)$ is finite.

In addition, two arrays of positive numbers a_{ij} and b_{ij} are given, where a_{ij} represents the loading-traveling time from i to j, and b_{ij} the unloading-traveling time from j to i.

The problem is to rearrange the numbers t_{ij}^k into s sequences such that

(2.1) each sequence is monotone increasing;

(2.2) if
$$t_{1}^{k_1} < t_{1}^{k_2}$$
 are consecutive numbers in any one of the s sequences, then

$$t_{i_2j_2}^{k_2} - t_{i_1j_1}^{k_1} \ge a_{i_1j_1} + b_{i_2j_1}$$
;

(2.3) s is minimal.

In other words, each sequence is a schedule for one tanker, and the objective is to meet the fixed schedule given by table t with a minimum number of tankers.

If, in the example, we take

$$a_{ij} = b_{ij} = \begin{vmatrix} 2 & 3 & 2 \\ 1 & 2 & 1 \end{vmatrix}$$

then a feasible schedule using seven tankers would be represented by the rearrangement

(1)
$$t_{11}^{1} = 1$$
, $t_{23}^{1} = 5$, $t_{22}^{1} = 7$, $t_{11}^{5} = 13$
(2) $t_{21}^{1} = 3$, $t_{21}^{2} = 6$, $t_{21}^{3} = 9$, $t_{12}^{2} = 15$
(3) $t_{11}^{2} = 4$, $t_{12}^{1} = 9$, $t_{22}^{4} = 15$
(4) $t_{13}^{1} = 6$, $t_{22}^{2} = 10$, $t_{23}^{3} = 15$
(5) $t_{11}^{3} = 7$, $t_{23}^{2} = 10$, $t_{22}^{3} = 13$
(6) $t_{13}^{2} = 12$
(7) $t_{11}^{4} = 10$.

3. REFORMULATION AS A PROGRAMMING PROBLEM

For convenience in exposition, we suppose that the numbers t_{ij}^k , a_{ij} , b_{ij} are positive integers. It will be clear that this is not essential to the method of solution.

First construct the table of sequences $T = (t_{1j}^k + a_{1j})$. Thus, T_{1j}^k are the times when tankers loaded at i will arrive at j. Define

 n_{Gi} = number of times $t_{ij}^k = \alpha$ occurs in row i of t; $n_{\beta j} = n$ number of times $T_{ij}^k = \beta$ occurs in column j of T; i.e., n_{Gi} is the number of tankers loading at i at time α and $n_{\beta j}$ is the number arriving at j at time β . Thus, n_{Gi} is defined for $\alpha = 1, 2, \ldots, \max_{j \in A} t_{ij}^k$ and $n_{\beta j}$ is defined for $\beta = 1, 2, \ldots, \max_{j \in A} T_{ij}^k$.

For any schedule, denote the number of reassignments from discharge point j at time β to loading point i at time α by Xaib.i. Then for all possible schedules, the inequalities

(3.1)
$$\sum_{\alpha,i} x_{\alpha i \beta j} \leq x_{\beta j}$$

$$\sum_{\beta,j} x_{\alpha i \beta j} \leq x_{\alpha i}$$
(3.2)
$$x_{\alpha i \beta j} \geq 0$$

are satisfied. In addition, it follows from (2.1) and (2.2) that

(3.3)
$$b_{ij} > \alpha - \beta$$
 implies $x_{\alpha i\beta j} = 0$.

The system (3.1) can be made into a system of equalities which

is formally of transportation type by introducing non-negative slack variables $x_{\alpha i}$, $y_{\beta j}$ and $z = \sum_{\alpha, i} \sum_{\beta, j} x_{\alpha i \beta j}$. Then (3.1) may be rewritten as

$$(3.4) \sum_{\alpha,1}^{\Sigma} x_{\alpha 1\beta j} + y_{\beta j} = x_{\beta j}, \quad y_{\beta j} \geq 0$$

$$\sum_{\beta,j}^{\Sigma} x_{\alpha 1\beta j} + x_{\alpha 1} = x_{\alpha 1}, \quad x_{\alpha 1} \geq 0$$

$$\sum_{\alpha,1}^{\Sigma} x_{\alpha 1} + z = \sum_{\alpha,1}^{\Sigma} x_{\alpha 1}, \quad z \geq 0$$

$$\sum_{\beta,j}^{\Sigma} y_{\beta j} + z = \sum_{\beta,j}^{\Sigma} x_{\beta j}, \quad z \geq 0$$

and hence each schedule leads to an integral solution of (3.2) and (3.4) which satisfies condition (3.3).

Conversely, given any integral solution of (3.2), (3.3), and (3.4), a schedule can be constructed from it as follows: Each $x_{\alpha i}$ will be the number of tankers which start their individual schedules at time α from loading point i; i.e., there will be $x_{\alpha i}$ sequences in the rearrongement which have a $t_{ij}^k = \alpha$ as first member. Delete one such $t_{ij}^k = \alpha$ from t; let $\beta_0 = T_{ij}^k = \alpha_0 + a_{ij}^k$. Since $N_{\beta_0 j_0} > 0$, at least one of the variables $x_{\alpha i} \beta_0 j_0 = \alpha_0 + a_{ij}^k \beta_0 j_0$ has a positive value. Select one such.

Case 1. $x_{\alpha_1 i_1 \beta_0 j_0} > 0$ was selected. Then there is a $t_{i_1 j_1}^{k_1} = \alpha_1$, since $n_{\alpha_1 i_1} > 0$. Assign α_1 as second member of the sequence. Observe that by (3.3), $\alpha_1 - \beta_0 \ge b_{i_1 j_0}$, hence

$$t_{i_1j_1}^{k_1} - t_{i_0j_0}^{k_0} = \alpha_1 - \beta_0 + a_{i_0j_0} \ge a_{i_0j_0} + b_{i_1j_0}$$

and (2.2) is satisfied. Strike out α_1 from t and reduce α_1 in β_0 , β_0 , α_1 by unity.

Case 2. $y_{0,0} > 0$ was selected. In this case the sequence ends with α_{0} . Reduce $y_{0,1}$, $N_{0,1}$ by unity.

with α_0 . Reduce $y_{\beta_0 j_0}$, $y_{\beta_0 j_0}$ by unity. If Case 1 obtained, let $\beta_1 = T_{i_1 j_1} = \alpha_1 + a_{i_1 j_1}$, and examine the values of the variables $x_{\alpha i \beta_1 j_1}$, $y_{\beta_1 j_1}$. Again one of these must be positive. Apply either Case 1 or Case 2 with α_1 playing the role of α_0 . Repetition of the procedure outlined must eventually end with the selection of some $y_{\beta_k j_k} > 0$ (since by (3.3), $\alpha \le \beta$ implies $x_{\alpha i \beta j} = 0$), thus completing one of the sequences. The others can be gotten in the same way.

Notice that while many schedules can be constructed from an integral solution of (3.2), (3.3), and (3.4), the only physical difference between two such is that there may be more than one tanker available at the same time at some discharge point, in which case they are interchangeable.

Thus, the tanker scheduling problem can be viewed as one of minimizing $\sum_{\mathbf{G},\mathbf{i}} \mathbf{x}_{\mathbf{G}\mathbf{i}}$, the number of sequences in a rearrangement (or what is the same thing, maximizing the variable z), over the set of integral solutions* of (3.2) and (3.4) in which the

^{*}Such whole number solutions always exist, e.g., take all $x_{\alpha i \beta j} = 0$. This corresponds to the worst possible schedule of assigning a different tanker for each trip.

variables designated by (3.3) are fixed at zero. But it is well known [1] that the maximum of a linear form defined over all solutions of (3.2) and (3.4) is always assumed at some integral solution, and it is easy to see that this fact is not altered by imposing additional constraints of the form $x_{aij} = 0$. Hence the scheduling problem can be solved by the simplex algorithm, since the nature of the algorithm is such as to obtain a required integral solution. Moreover, the algorithm is extremely simple to apply when the problem is of transportation type, as is this one.

It is obvious also that linear programming can be used to optimize schedules with respect to other costs. For example, it would be simply a matter of changing the minimizing form, holding z fixed, to find a schedule for a given number of tankers which has the least sailing time.

4. A NUMERICAL EXAMPLE

We continue with the example of $\S 2$. First form the table of arrival times

imes 1 2 3
$$T = \begin{bmatrix} 1 & 3 & 6 & 9 & 12 & 15 & 12 & 18 & 8 & 14 \\ 2 & 4 & 7 & 10 & 13 & 9 & 12 & 15 & 17 & 6 & 11 & 16 \end{bmatrix}$$

Using this table and the one of loading times, compute all n

and $N_{\beta j}$. After discarding those rows and columns having $n_{\alpha j}$ or $N_{\beta j}$ zero, one is left with the transportation problem whose constraints are indicated schematically in Fig. 1. Crossed out cells mean that the corresponding variable is constrained to be zero by (3.3). The solution shown in Fig. 1 corresponds to the schedule using seven tankers given in §2. This is a degenerate solution to the programming problem and so it is necessary in applying the simplex algorithm to pick out other basic variables having zero values. One way of doing this is shown in Fig. 1 by the placement of the O's.

An optimal solution, reached after a few iterations, is shown in Fig. 2. It corresponds to the following six-tanker schedule:

$t_{11}^1 = 1$,	$t_{23}^1 = 5$,	$t_{22}^1 = 7$,	t ⁵ ₁₁ = 13
t ₂₁ = 3,	$t_{21}^2 = 6$,	$t_{21}^3 = 9$,	$t_{13}^2 = 12$
t ² ₁₁ = 4,,	$t_{12}^1 = 9$,	$t_{22}^{4} = 15$	
$t_{11}^3 = 7$,	$t_{23}^2 = 10,$	$t_{21}^{4} = 12,$	$t_{12}^2 = 15$
t ⁴ ₁₁ = 10,	$t_{22}^3 = 13$	•	
	$t_{21}^{1} = 3,$ $t_{11}^{2} = 4,,$ $t_{13}^{1} = 6,$ $t_{11}^{3} = 7,$	$t_{21}^{1} = 3$, $t_{21}^{2} = 6$, $t_{11}^{2} = 4$, $t_{12}^{1} = 9$, $t_{13}^{1} = 6$, $t_{22}^{2} = 10$, $t_{11}^{3} = 7$, $t_{23}^{2} = 10$,	$t_{11}^{1} = 1 , t_{23}^{1} = 5 , t_{22}^{1} = 7 ,$ $t_{21}^{1} = 3 , t_{21}^{2} = 6 , t_{21}^{3} = 9 ,$ $t_{11}^{2} = 4 , t_{12}^{1} = 9 , t_{22}^{4} = 15 $ $t_{13}^{1} = 6 , t_{22}^{2} = 10 , t_{23}^{3} = 15 $ $t_{11}^{3} = 7 , t_{23}^{2} = 10 , t_{21}^{4} = 12 ,$ $t_{11}^{4} = 10 , t_{22}^{3} = 13 .$

100 - AEUFFEL & 18577 FO .

7 8 9

1 2

10 11 12

1

13 14 15

16 17 18

3

6

1 2

RM-1328

n_{a1}

REFERENCES

- 1. G. Dantzig, "Application of the Simplex Method to a Transportation Problem," Activity Analysis of Production and Allocation, Cowles Commission Monograph No. 13, 1951.
- 2. J. Robinson and J. Walsh, Routing of Empties for Fixed—Schedule Transportation, The RAND Corporation, Memorandum M-406, June 12, 1950.
- 3. C. Tompkins, Discrete Problems and Computers, INA-S3-5, November 17, 1952, pp. 14-16.